

AD \_\_\_\_\_

Award Number: DAMD17-97-1-7266

TITLE: The Role of NF-kB in Normal and Transformed Mammary  
Epithelium

PRINCIPAL INVESTIGATOR: Fiona Yull, Ph.D.  
Dr. Kerr

CONTRACTING ORGANIZATION: Vanderbilt University Medical Center  
Nashville, Tennessee 37232

REPORT DATE: August 2000

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

**REPORT DOCUMENTATION PAGE**Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

<b>1. AGENCY USE ONLY (Leave blank)</b>		<b>2. REPORT DATE</b> August 2000	<b>3. REPORT TYPE AND DATES COVERED</b> Final (1 Aug 97 - 31 Jul 00)	
<b>4. TITLE AND SUBTITLE</b> The Role of NF- $\kappa$ B in Normal and Transformed Mammary Epithelium			<b>5. FUNDING NUMBERS</b> DAMD17-97-1-7266	
<b>6. AUTHOR(S)</b> Fiona Yull, Ph.D. Dr. Kerr			<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Vanderbilt University Medical Center Nashville, Tennessee 37232  E-MAIL: Fiona.yull@mcmail.vanderbilt.edu Department of Microbiology and Immunology				
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b>  U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			<b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>				
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for public release; distribution unlimited			<b>12b. DISTRIBUTION CODE</b>	
<b>13. ABSTRACT (Maximum 200 Words)</b>  A knowledge of the molecular signaling pathways which, when perturbed, result in mammary cell transformation would provide vital information in the fight against breast cancer. The NF- $\kappa$ B family of transcription factors are a major component of pathways regulating cell proliferation, differentiation and death. We are attempting to determine their role in the development of normal and transformed mammary epithelium. Our hypothesis is that expression of specific NF- $\kappa$ B factors is necessary for the proper differentiation and maturation of mammary epithelium and that altered expression of $\kappa$ B factors can effect transformation of mammary epithelium. We have created lines of transgenic mice which overexpress a family member suspected to be involved in carcinogenesis (NF $\kappa$ B2) or a dominant inhibitor of NF- $\kappa$ B activity (I $\kappa$ B $\alpha$ - $\Delta$ N) in the mammary gland. These murine models are now being analyzed to determine how proliferation, differentiation and death are changed in response to altered $\kappa$ B activity.				
<b>14. SUBJECT TERMS</b> Breast Cancer, NF- $\kappa$ B, transgenics			<b>15. NUMBER OF PAGES</b> 34	
			<b>16. PRICE CODE</b>	
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified	<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified	<b>20. LIMITATION OF ABSTRACT</b> Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)  
Prescribed by ANSI Std. Z39-18  
298-102

## FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

\_\_\_\_ Where copyrighted material is quoted, permission has been obtained to use such material.

\_\_\_\_ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

KEY Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

KEY In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

\_\_\_\_ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

KEY In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

KEY In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

\_\_\_\_ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Foran E. Tull

PI - Signature

2/31/00

Date

## Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	4
Introduction.....	5
Body.....	5
Key Research Accomplishments.....	9
Reportable Outcomes.....	10
Conclusions.....	11
References.....	12
Appendices.....	14

## INTRODUCTION

The nuclear factor-kappaB, NF- $\kappa$ B, family of transcription factors play roles in normal functions such as cellular proliferation, differentiation, migration, and programmed cell death (1, 2). They are also implicated in the pathogenesis of cancer, particularly in cells of the immune system (3). However, little is known concerning the precise mechanisms by which this occurs. Our research has two major goals. The first is to characterize the role of the transcription factor complex in normal mammary development. The second is to investigate the reported correlation between the presence of an unusual isoform of NF- $\kappa$ B, NF $\kappa$ B2, and breast adenocarcinoma (4). Our strategy has been to generate novel transgenic models. In the first model, the activity of NF- $\kappa$ B is blocked in mammary epithelial cells by the expression of a dominant inhibitor (IkB $\Delta$ N) to determine the effects of disturbing normal signalling pathways. In the second model, NF $\kappa$ B2 (p100) is over-expressed in mammary epithelial cells to determine whether this protein can induce cellular transformation. If this were the case we would have generated a novel murine model of mammary tumorigenesis. In both cases we have used the promoter of the major whey protein of ruminants,  $\beta$ -lactoglobulin (BLG) to target transgene expression. We have generated four lines of transgenic mice targeting IkB $\Delta$ N to the mammary gland, designated MAN for mammary - $\alpha$  $\Delta$ N. We have also produced four lines of mice targeting overexpression of NF $\kappa$ B2 (p100), designated BIN for BLG-inhibitor. The lines are currently being characterized.

## BODY

The Rel/NF- $\kappa$ B family of transcription factors represents a distinct subset of nuclear transactivating factors whose activity is inducible via control of its nuclear localization. These enhancer binding proteins are sequestered in the cytoplasm by inhibitory molecules, termed IkBs. Upon stimulation the IkB is degraded, and the  $\kappa$ B factors are released to enter the nucleus and associate with their cognate DNA binding sites initiating gene transcription (5). Authentic NF- $\kappa$ B is a heterodimer consisting of a 50 kD polypeptide (NF $\kappa$ B1=p50) (6) and a 65 kD polypeptide (RelA=p65)(7) whose primary sequence reveals that the amino-terminal halves of both p50 and p65 are highly homologous to that of the *rel* protein (defining the Rel homology domain, RHD). The RHD encompasses the DNA-binding, dimerization, IkB-binding, basal transcription factor binding, and nuclear localizing domains for these proteins. The carboxy terminal domains of RelA(p65) and cRel contain strong transcriptional transactivation regions (8). In contrast, p50 homodimers bind DNA but are thought to block transcription. In vertebrates, other genes also encode factors which participate in the NF- $\kappa$ B complex and bind to  $\kappa$ B enhancer elements: NF $\kappa$ B2 (p52 and its precursor p100)(9), relb (RelB), and v-rel.

IkB $\alpha$  forms a complex with heterodimeric NF- $\kappa$ B and inhibits the DNA binding activity (10). Following stimulation with a wide variety of distinct agents, the IkB- $\alpha$  protein is phosphorylated on serine residues in the amino-terminus targeting it for destruction by the ubiquitination/proteasome (26S) degradation pathway, allowing the translocation of NF- $\kappa$ B to the nucleus (10,11,12). A mutation in the IkB- $\alpha$  which removes these serine residues creates an IkB- $\alpha$  protein (termed IkB $\Delta$ N) behaves as a transdominant inhibitor (13). This has provided an effective means by which to "capture" NF- $\kappa$ B in a cytoplasmic-bound state, thereby preventing its transactivating capabilities (14).

NF $\kappa$ B2 was originally identified via its involvement in a lymphoma-associated chromosomal translocation and is expressed as a 3.2 kilobase mRNA encoding a 100 kDa protein (15). Following proteolytic cleavage to the mature p52 kDa protein, NF $\kappa$ B2-p52 can heterodimerize with all known  $\kappa$ B factors and activate the transcription of genes containing  $\kappa$ B-enhancer elements. The NF $\kappa$ B2-p100 precursor has activity on its own; p100 behaves as an IkB

molecule by sequestering NF- $\kappa$ B complexes in the cytoplasm. In one lymphoid-tumor case reported, overexpressed NF $\kappa$ B2 proteins resulted in abnormal transcriptional regulatory properties (16) suggesting that alteration in NF $\kappa$ B2-regulated gene expression may be causal for transformation. These results suggest that overexpression of either the precursor molecule, p100, or its processed fragment, p52, might alter the normal cellular NF- $\kappa$ B complexes and disregulate cellular homeostasis.

In order to target expression of our transgenes to mammary epithelial cells we have used the promoter of the major whey protein of ruminants,  $\beta$ -lactoglobulin (BLG). Despite the fact that rodents do not have an endogenous BLG gene transgenic mice carrying BLG constructs express specifically and abundantly in the murine mammary gland in an appropriate temporal pattern (17,18). Transgenes using 5' and 3' BLG flanking sequences can target expression of heterologous genes in an appropriate developmental pattern to the mammary (19,20,21). In the previous annual report we had succeeded in generating four lines of transgenic mice targeting I $\kappa$ B $\Delta$ N to the mammary gland, designated MAN for mammary - $\alpha$ \DeltaN. We had also produced four lines of mice targeting overexpression of NF $\kappa$ B2 (p100), designated BIN for BLG-inhibitor. This report details further characterization of these lines.

### **Results:**

Specific aim 1 (Task 1) for this proposal was to define the expression pattern of NF- $\kappa$ B factors in normal murine mammary gland development. This aim was addressed in collaboration with Dana Brantley (awarded a pre-doctoral fellowship - Award Number DAMD17-97-1-7017) and has led to publication of a manuscript (22). See relevant final report for details.

Specific aim 2 proposed the generation and characterization of two types of transgenics targeting the transdominant inhibitor I $\kappa$ B $\Delta$ N or NF $\kappa$ B2 expression to the murine mammary gland. Four lines representing each type of transgenic have been generated (Tasks 2- 5). All lines have been backcrossed five generations onto an FVB strain background for future carcinogenesis studies. A repository of samples has been collected representing a range of developmental stages throughout pregnancy, lactation and regression (Tasks 6-7). These include; virgin, 10.5 and 16.5 dpc (days post coitem, ie. pregnant), 1.5 and 9.5 lactating and 1, 3, 5 and 40 days and 5 weeks post forced wean (at day 10 of lactation) and greater than 1 year. At each stage one gland is removed for histological processing, one is processed in TriZol reagent for RNA extraction, one is flash frozen for isolation of cytoplasmic and nuclear proteins for EMSA, and one gland is preserved for whole mount gland preparation. The particular gland collected for each type of analysis remains constant. Collection of these stages represents a major time commitment. Transgenic animals are identified by tail biopsy and PCR. Female transgenics are set up in timed matings and day post coitem noted by detection of a vaginal plug. Then animals are sacrificed on the relevant day of pregnancy, lactation etc. Breeding studies to generate BLG- $\Delta$ N/HLL:MAN and BLG- $\Delta$ N/HLL:BIN double transgenics are well underway and preliminary analysis completed (Tasks 8-9). The cross breeding to generate BLG-NF $\kappa$ B2/p53-/- lines has not been started. The p53-/- carcinogenesis model mice are on the FVB strain background and it was decided that it would be prudent to backcross the MAN and BIN transgenics onto the same genetic background before commencing these studies. Such backcrosses were not included in the original Statement of Work and require considerable time investment.

To summarize in more detail progress to date:

Primary characterization of expression has been completed by Northern blot analysis. Expression of I $\kappa$ B $\Delta$ N is detected in all four separately derived MAN transgenic lines ie. MAN8, 13, 18 and 19. Expression of p100 has also been detected in three out of four separately derived

NF- $\kappa$ B2 (p100) lines ie. BIN10, 16 and 18. Offspring from the fourth founder animal, representing the BIN6 line, whilst carrying the transgene have either extremely low expression or a "silent transgene" (see prev. reports). We have elected to continue sample collection and backcrossing with the BIN6 line as it has proved to be a difficult line to propagate suggestive of some transgene effect.

Northern analysis included investigation of  $\beta$ -casein expression. Although the transgenics appear capable of nursing litters we were interested in whether transgene expression alters milk protein production. We have analysed total RNA samples representing 16.5 dpc, 1.5 days lactation and 9.5 days lactation in all four I $\kappa$ B $\Delta$ N expressing (MAN) lines and in all four BIN lines. In all lines tested and at all stages, the expression of either transgene does not appear to have a significant effect on  $\beta$ -casein expression.

In order to determine whether transgene expression perturbs normal NF- $\kappa$ B activity, we have analyzed the binding activity of protein extracts by electrophoretic mobility shift assay (EMSA) using a probe derived from the HIV-LTR that represents a consensus binding site for NF- $\kappa$ B. We have investigated binding activity at the 16.5 dpc and 9.5 lactating points. The four p100 (BIN) lines have so far shown no significant differences at the 16.5 dpc stage. However, analysis of the 9.5 lactation stage detects intriguing differences (see previous report). In non-transgenic animals there is barely detectable binding activity during lactation (22). In transgenics a strong shifted complex is present. This complex can be competed for by competitor oligos. Supershift experiments are being attempted using anti-p52, anti-p65 and anti-p50. Whilst no supershifts indicating the presence of p52 or p65 are detected we have been experiencing some technical difficulties with these antibodies such that further investigation is underway to confirm this result. It appears as though the p50 protein represents a large component of the transgenic complex. Preliminary analysis of protein extracts representing the 9.5 day lactation stage of I $\kappa$ B $\Delta$ N (MAN) transgenics has detected no significant alteration in NF- $\kappa$ B binding activity with no retarded complexes in transgenics or controls. However, EMSA of protein samples from the 16.5 dpc stage show subtle changes. At this stage of pregnancy in a non-transgenic animal there is strong NF- $\kappa$ B binding activity (22). Formation of this complex is inhibited by unlabelled oligonucleotide. There appears to be a decrease in the retardation of the complex (or a decrease in the presence of the upper band) in separately derived transgenic lines MAN8 and MAN19. This is interesting in the light of our observation of a decrease in cyclinD1 RNA expression (as determined by RPA analysis) in these lines.

One of the possible predicted effects of perturbation of NF- $\kappa$ B activity would be to alter proliferation. Intraperitoneal injections of BrdU were administered to animals at day 16.5 of pregnancy. Mice were sacrificed 4 hours later, mammary glands fixed overnight in 4% paraformaldehyde, dehydrated and paraffin embedded sections prepared. Sections were processed using a BrdU staining kit (Zymed) to investigate proliferation of the alveolar epithelial cells (see previous report). Proliferating cells were detected in all samples tested with no obvious qualitative differences in proliferation between transgenics and non-transgenic controls. However, some animals exhibited histological differences including a relative decrease in the proportion of alveolar cells populating the "web-like" stroma, a less uniform morphology and an increase in the number of ductal structures. A limited number of samples have been tested. We plan to expand this set and carry out more quantitative measurements by counting relative numbers of proliferating cells. Particularly in light of the results from the work supported by pre-doctoral fellowship - Award Number DAMD17-97-1-7017 in this laboratory which suggest that in the absence of a major inhibitor of NF- $\kappa$ B activity the developing mammary gland becomes hyperplastic.

A continuing concern is the potential for between animal variation (especially as the original mice are B6/D2 (ie. a mixture of both the C57Bl6 and DBA mouse strains). At this stage we are still unable to confirm whether there may be a measure of mosaicism in the expression of the

transgenes. We attempted to address this issue using immunohistochemistry of collected paraffin sections, however, our attempts were unsuccessful due to technical difficulties with the available commercial p52 antibody. We immunized rabbits in an effort to produce a COOH-terminal I $\kappa$ B- $\alpha$  antibody which would be effective for both immunohistochemical and western analyses of the I $\kappa$ B $\Delta$ N (MAN) transgenics. A second paired set of immunizations were to produce N- and COOH-terminal antibodies to the p100 protein. These would enable us to effect immunohistochemical and western studies on the BIN lines and to distinguish between the unprocessed p100 form of the protein and the truncated p52 protein. The terminal bleeds of the rabbits have been completed. An N-terminal antibody (antigen designed and injected with the original set) has proved to be very useful for other studies in the laboratory. However, this antibody cannot be used to recognize the N-terminally deleted form of I $\kappa$ B $\Delta$ N used in our transgenics. The COOH-terminal antibody has not proved to be useful as it cross reacts with endogenous mouse MAD3. The p100 sera remain to be tested.

Mammary epithelial transformation and neoplastic progression, like most cancers, involves cooperative changes in more than one oncogenic or tumor-suppressor pathway. Eventually, it is our intention to cross selected BIN and MAN lines with mice expressing the mouse mammary tumor virus (MMTV) driven-activated *neu* transgene, which reproducibly develop clonal mammary tumors, to determine whether transgene expression can either inhibit or cooperate in producing mammary tumorigenesis. Furthermore, it would be interesting to breed BLG-NF $\kappa$ B2/p100 expressing mice with mice expressing the most common p53 mutation (Arg<sup>172</sup>-His) in human breast cancers to explore the ability of NF $\kappa$ B2 to act as a dominant oncogene in the absence of the well characterized tumor suppressor, p53. These interbreedings would explore the cooperative nature between these NF- $\kappa$ B components and known mammary carcinogenic signal transduction and genetic pathways in the hopes of better understanding the mechanistic interactions of the oncogenic pathways in human mammary tumorigenesis. The lines of mice described above exist on an FVB strain background whereas our lines were generated on a B6/D2 background. In order to avoid the potential difficulties of interpreting data on a background consisting of a genetic mix of three strains we have backcrossed our lines onto the FVB strain. We have achieved at least 5th generation backcross and are characterizing these animals for transgenic phenotype.

We have generated double transgenics between our mammary lines and reporter lines generated in this lab designated HLL (23). The HIV-1 LTR has been extensively characterized as a NF- $\kappa$ B responsive promoter (24). These lines carry the NF- $\kappa$ B responsive human immunodeficiency virus-1 long terminal repeats, HIV-LTR, promoter fused to luciferase which enables direct analysis of the expression of an *in vivo* NF- $\kappa$ B reporter. This provides an extremely valuable system to determine whether expression of I $\kappa$ B $\Delta$ N is blocking NF- $\kappa$ B activity *in vivo*, and to determine whether the overexpression of NF $\kappa$ B2 results in an active transcription factor or whether the full length protein (p100) is behaving as an inhibitor of other NF- $\kappa$ B complexes and thus, preventing transcriptional activation. Our results so far are as follows:

Preliminary analysis of double transgenic animals for luciferase expression at 16.5dpc stated as relative light units per  $\mu$ g total protein (rlu/ $\mu$ g).

I $\kappa$ B $\alpha$ - $\Delta$ N (MAN) transgenics have relatively decreased NF- $\kappa$ B activity.

HLL	1106 $\pm$ 288 rlu/ $\mu$ g
HLL/MAN	537 $\pm$ 268 rlu/ $\mu$ g

p100-NF $\kappa$ B2 (BIN) transgenics have relatively increased NF- $\kappa$ B activity.

HLL	24 $\pm$ 8 rlu/ $\mu$ g
HLL/BIN10	118 $\pm$ 79 rlu/ $\mu$ g
HLL/BIN16	73 $\pm$ 8 rlu/ $\mu$ g



[The difference in control single HLL transgenic background between the MAN study and the BIN study is the result of use of a different protocol and type of luminometer with decreased background levels of luciferase activity]

The FVB strain background is believed to be more susceptible to tumor formation than the current B6/D2 strain background. In addition very limited effects of leaky transgene expression would be expected in virgin animals (as the promoter used normally directs expression of a milk protein). Thus we were surprised on examining haematoxylin and eosin stained sections from older, virgin BIN transgenic animals to note some morphological differences relative to control glands ( see previous report). This may be an indication of future more profound effects on older animals on the FVB strain background after a couple of rounds of pregnancy. We are collecting samples from older FVB transgenic virgins and have set up a small pilot study of control and BIN transgenics on a continuous mating schedule.

### **Key Research Accomplishments**

- 1) Generation of four lines of transgenic mice expressing a dominant inhibitor of NF $\kappa$ B (IkB $\Delta$ N).
- 2) Generation of four lines of mice carrying a mammary specific NF $\kappa$ B2 (p100) transgene.
- 3) Characterization of above lines confirming mammary specific transgene expression.
- 4) Transgene expression has no profound effect on milk production.
- 5) Backcross of 8 lines to 5th generation onto the FVB strain background for future carcinogenesis studies.
- 6) Putative generation of antibodies to the COOH terminal of IkB $\alpha$  and both termini of NF $\kappa$ B2.

## Reportable Outcomes

### 1) Manuscript:

Brantley, D.M., Yull, F.E., Muraoka, R.S., Hicks, D., Cook, C.M. and Kerr, L.D. Dynamic expression and activity of NF- $\kappa$ B during post-natal mammary gland morphogenesis. (In Press, Mechanisms of Development).

### Abstracts:

Yull, F.E., Brantley, D.M., Seetharaman, R., Bradberry, J. and Kerr, L.D. Transgenic models to study the role of NF- $\kappa$ B in mammary morphogenesis and neoplasia. Era of Hope, Department of Defense Breast Cancer Research Program Meeting, Atlanta (2000).

Brantley, D.M., Chen, C-L., Yull, F.E. and Kerr, L.D. Nuclear factor-kappa B (NF- $\kappa$ B) factors in mammary epithelial development and proliferation. Era of Hope, Department of Defense Breast Cancer Research Program Meeting, Atlanta (2000).

Yull, F.E., Brantley, D.M., and Kerr, L.D. Transgenic models to investigate the role of NF- $\kappa$ B in normal and transformed mammary epithelium. Society for Developmental Biology SE Regional Meeting, Emory University, (1998).

2) Accumulation of major repository of tissue, RNA, protein, whole mount and paraffin section samples representing multiple stages of mammary development from wild-type and 8 transgenic lines. Samples collected from both the original mixed C57B16/DBA animals and 5th generation backcross to the FVB strain.

3) The I $\kappa$ B $\Delta$ N transgenics (MAN) represent a novel model which will be exploited to investigate the roles of NF $\kappa$ B in normal mammary development. These will permit examination of the role of NF $\kappa$ B in pregnancy (proliferation/differentiation), lactation (secretion) and post-weaning (apoptosis).

4) The NF $\kappa$ B2 transgenics (BIN) represent a novel model which will be exploited to investigate the role of NF- $\kappa$ B in carcinogenesis.

## CONCLUSIONS

Our hypothesis was that the nuclear expression of specific NF- $\kappa$ B factors is necessary for the proper differentiation and maturation of mammary epithelium and that alteration in the expression of  $\kappa$ B factors can effect transformation of mammary epithelium. We have established lines of transgenic mice in which NF- $\kappa$ B levels are perturbed by overexpression of an NF- $\kappa$ B family member correlated with human mammary carcinogenesis (NF $\kappa$ B2) or by expression of a dominant negative factor (I $\kappa$ B $\alpha$ - $\Delta$ N), inhibitory towards nuclear NF- $\kappa$ B activity, in the mammary gland. These transgenics are designated BIN and MAN respectively. Analysis of the transgenic lines is continuing. Indications are that the transgenes are able to perturb NF- $\kappa$ B activity at different stages but further studies are required to determine the nature of these alterations. No profound effect on the ability of either transgene to alter milk protein production has been detected. Suspected effects on proliferation are proving difficult to confirm.

We have backcrossed our animals onto the FVB strain to alleviate any potential difficulties associated with variation due to the original C57Bl6/DBA mixed genetic background and are continuing characterizations on the FVB genetic background. The FVB backcrosses are also required for tumorigenesis studies which we hope to secure funding to continue. We have also attempted to generate antibodies to determine whether the transgenes are expressed consistently across entire glands or whether some degree of mosaic expression may be present.

We have utilized transgenic lines generated in this laboratory in which the NF- $\kappa$ B responsive HIV-LTR directs expression of luciferase, enabling direct *in vivo* analysis of NF- $\kappa$ B activity. These transgenic lines have been crossed with the mammary transgenics. Results suggest that expression of the dominant inhibitor of NF- $\kappa$ B, I $\kappa$ B- $\alpha$  $\Delta$ N, blocks  $\kappa$ B activity *in vivo*. Overexpression of NF $\kappa$ B2 appears to result in an active transcription complex during pregnancy. Initial histological studies also suggest that expression of the transgenes can result in aberrant morphogenesis of the mammary gland during pregnancy.

At this time no clear indications of transformation are apparent in the BIN lines. This may be due, in part to genetic susceptibility. Representative lines have been backcrossed five generations onto the FVB genetic background, known to be susceptible to tumorigenesis for further studies. These are scheduled to include generation of MMTV-neu/BIN double transgenics to investigate the potential for co-operative effects.

At this stage indications are that the transgenes differentially affect NF $\kappa$ B complex formation and activity. More information is necessary before it can be determined whether dysregulation of NF- $\kappa$ B complex formation results in altered regulation of gene products leading to unscheduled growth, misregulated differentiation, altered rates of apoptosis, excess angiogenesis or invasiveness.

In summary, in relation to the original statement of work, samples collected for technical objective 1 have been included in the results reported for award #DAMD17-97-1-7017. Tasks 2-7 of technical objective 2 have been completed. Task 8 has been completed for the breeding studies to generate BLG- $\Delta$ N/HLL animals and characterization as per Task 9 is underway.

## REFERENCES

1. Verma, I.M., Stevenson, J.K., Schwarz, E.M., Van Antwerp, D., and Miyamoto, S. 1995. Rel/NF- $\kappa$ B/I $\kappa$ B family: intimate tales of association and dissociation. *Genes and Devel.* 9:2723-2735.
2. Kerr, L.D. and Verma, I.M. 1992. Signal transduction: The nuclear target. *Curr. Opin. Cell Biol.* 4: 496-501.
3. Gilmore, T.D., Koedood, M., Piffat, K.A. And White, D.W. 1996. Rel/NF- $\kappa$ B/I $\kappa$ B proteins and cancer. *Oncogene* 13: 1367-1378.
4. Dejardin, E., Bonizzi, G., Bellahcene, A., Castronovo, V., Merville, M.-P., and Bours, V. 1995. Highly-expressed p100/p52 (NF $\kappa$ B2) sequesters other NF- $\kappa$ B-related proteins in the cytoplasm of human breast cancer cells. *Oncogene* 11: 1835-1841.
5. Nolan, G.P., Baltimore, D. 1992. The inhibitory ankyrin and activator rel proteins. *Curr. Opin. Genet. Dev.* 2: 211-220.
6. Ghosh, S., Gifford, A.M., Riviere, L.R., Tempst, P., Nolan, G.P., and Baltimore, D. 1990. Cloning of the p50 DNA binding subunit of NF- $\kappa$ B: Homology to *rel* and *dorsal*. *Cell* 62: 1019-1029.
7. Nolan, G.P., Ghosh, S., Liou, H.-C., Tempst, P., Baltimore, D. 1991. DNA binding and I $\kappa$ B inhibition of the cloned p65 subunit of NF- $\kappa$ B, a rel-related polypeptide. *Cell* 64: 961-969.
8. Bull, P. Morley, K.L., Hoekstra, M.F., Hunter, T., Verma, I.M. 1990. The mouse *c-rel* protein has an N-terminal regulatory domain and a C-terminal transcriptional transactivation domain. *Mol. Cell. Biol.* 10: 5473-5485.
9. Scheinman, R.I., Beg, A.A., and Baldwin, A.S. 1993. NF- $\kappa$ B p100 (Lyt-10) is a component of HSTF1 and can function as an I $\kappa$ B-like molecule. *Mol. Cell. Biol.* 13: 6089-6101.
10. Brockman, J.A., Scherer, D.C., McKinsey, T.A., Hall, S.M., Qi, X., Lee, W.Y., Ballard, D.W. 1995. Coupling of a signal response domain in I $\kappa$ B $\alpha$  to multiple pathways for NF- $\kappa$ B activation. *Mol. Cell. Biol.* 15: 2809-2818.
11. Treanckner, E. B.-M., Pahl, H.L., Henkel, T., Schmidt, K.N., Wilk, S., Baeuerle, P.A. 1995. Phosphorylation of human I $\kappa$ B- $\alpha$  on serines 32 and 36 controls I $\kappa$ B- $\alpha$  proteolysis and NF- $\kappa$ B activation in response to diverse stimuli. *EMBO J.* 14: 2876-2883.
12. Chen, C-L, Yull, F.E. And Kerr, L.D. 1999. Differential serine phosphorylation regulates I $\kappa$ B- $\alpha$  inactivation. *Biochem. Biophys. Res. Comm.* 257: 798-806.
13. Inoue, J.-i., Kerr, L.D., Rashid, D., Davis, N., Bose, H.R., Verma, I.M. 1992. Direct association of pp40/I $\kappa$ B $\beta$  with rel/NF- $\kappa$ B transcription factors: Role of ankyrin repeats in the inhibition of DNA binding activity. *Proc. Natl. Acad. Sci USA* 89: 4333-4337.
14. Bushdid, P.B., Brantley, D.M., Yull, F.E., Blaeuer, G.L., Hoffman, L.H., Niswander, L. and Kerr, L.D. 1998. Inhibition of NF- $\kappa$ B activity results in disruption of the apical ectodermal ridge and aberrant limb morphogenesis. *Nature* 392: 615-618.
15. Mercurio, F., DiDonato, J., Rosette, C., and Karin, M. 1992. Molecular cloning and characterization of a novel Rel/NF- $\kappa$ B family member displaying structural and functional homology to NF- $\kappa$ B/p50/p105. *DNA Cell Biol.* 11: 523-537.
16. Zang, J., Chang, C.-C., Lombardi, L., and Dalla-Favera, R. 1994. Rearranged NF $\kappa$ B2 gene in the HUT78 T-lymphoma cell line codes for a constitutively nuclear factor lacking transcriptional repressor functions. *Oncogene* 10: 1931-1937.
17. Simons, J.P., McClenaghan, M. and Clark, A.J. 1987. Alteration of the quality of milk by expression of sheep B-lactoglobulin in transgenic mice. *Nature* 328: 530-532.
18. Harris, S., McClenaghan, M., Simons, J.P., Ali, S. and Clark, A.J. 1991. Developmental

- regulation of the sheep beta-lactoglobulin gene in the mammary gland of transgenic mice. *Developmental Genetics* **12**: 299-307.
19. Archibald, A.L., McClenaghan, M., Hornsey, V., Simons, J.P. and Clark, A.J. 1990. High level expression of biologically active human  $\alpha$ 1-antitrypsin in transgenic mice. *Proc. Natl. Acad. Sci. USA* **87**: 5178-5182.
  20. Whitelaw, C.B.A., Harris, S., McClenaghan, M., Simons, J., and Clark, A.J. 1992. Position-independent expression of the ovine  $\beta$ -lactoglobulin gene in transgenic mice. *Biochem. J.* **286**: 31-39.
  21. Yull, F., Harold, G., Wallace, R., Cowper, A., Percy, J., Cottingham, I., and Clark, A.J. 1995. Fixing human factor IX (fIX): Correction of a cryptic RNA splice enables the production of biologically active fIX in the mammary gland of transgenic mice. *Proc. Natl. Acad. Sci. USA* **92**: 10899-10903.
  22. Brantley, D.M., Yull, F.E., Muraoka, R.S., Hicks, D., Cook, C.M. and Kerr, L.D. Dynamic expression and activity of NF- $\kappa$ B during post-natal mammary gland morphogenesis. (In Press, *Mechanisms of Development*).
  23. Yull, F.E., Blackwell, T.S., Chen, C-L, Venkatakrishnan, A., Blackwell, T.R., Hicks, D.J., Lancaster, L.H., Christman, J.W. and Kerr, L.D. 1998. NF- $\kappa$ B activation and cytokine production in a transgenic mouse model of endotoxin-induced lung inflammation. (In Press, *Am. J. Crit. Care Med.*).
  24. Kretzschmar, M., Meisterernst, M., Scheidereit, C., Li, G., Roeder, R.G. 1992. Transcriptional regulation of the HIV-1 promoter by NF- $\kappa$ B. *Genes. & Devel.* **6**:761-774.

# Appendices

## Bibliography

### Publications:

Brantley, D.M., Yull, F.E., Muraoka, R.S., Hicks, D., Cook, C.M. and Kerr, L.D. Dynamic expression and activity of NF- $\kappa$ B during post-natal mammary gland morphogenesis. (In Press, Mechanisms of Development).

### Meeting Abstracts:

Yull, F.E., Brantley, D.M., Seetharaman, R., Bradberry, J. and Kerr, L.D. Transgenic models to study the role of NF- $\kappa$ B in mammary morphogenesis and neoplasia. Era of Hope, Department of Defense Breast Cancer Research Program Meeting, Atlanta (2000).

Brantley, D.M., Chen, C-L., Yull, F.E. and Kerr, L.D. Nuclear factor-kappa B (NF- $\kappa$ B) factors in mammary epithelial development and proliferation. Era of Hope, Department of Defense Breast Cancer Research Program Meeting, Atlanta (2000).

Yull, F.E., Brantley, D.M., and Kerr, L.D. Transgenic models to investigate the role of NF- $\kappa$ B in normal and transformed mammary epithelium. Society for Developmental Biology SE Regional Meeting, Emory University, (1998).

### Personel:

Partial salary support for Fiona E. Yull, D.Phil., Research Assistant Professor.

## GENE EXPRESSION PATTERNS SECTION

### Dynamic Expression and Activity of NF- $\kappa$ B During Post-Natal Mammary Gland Morphogenesis

Dana M. Brantley<sup>1</sup>, Rebecca S. Muraoka<sup>5</sup>, Fiona E. Yull<sup>1</sup>, Donna J. Hicks<sup>1</sup>, Christopher M. Cook<sup>2</sup>, & Lawrence D. Kerr<sup>1,2,3,4\*</sup>

Departments of Microbiology & Immunology<sup>2</sup>, Cell Biology<sup>1</sup>, and Hematology and Oncology<sup>5</sup>, Vanderbilt University School of Medicine and the Vanderbilt-Ingram Cancer Center<sup>3</sup>, 1161 21st Avenue South, Nashville, TN 37232-2363, USA; Immunogenetics Section, Division of Allergy, Immunology and Transplantation (DAIT), National Institute of Allergy and Infectious Diseases (NIAID), National Institutes of Health (NIH), Bethesda, MD 20892, USA<sup>4</sup>

Key Words: NF- $\kappa$ B, p105, p50, RelA, I $\kappa$ B $\alpha$ , mammary, epithelium, proliferation, differentiation, involution, apoptosis, transgenic

\*Lawrence D. Kerr, Ph.D.  
Vanderbilt University School of Medicine  
A-4314 MCN  
1161 21st Avenue South  
Nashville, TN 37232-2363  
(615) 343-2568  
(615) 343-2569 FAX  
Lkerr@niaid.nih.gov

CORRESPONDENCE and requests for materials should be addressed to Lawrence D. Kerr

## GENE EXPRESSION PATTERNS SECTION

### Abstract

The Rel/NF- $\kappa$ B family of transcription factors has been implicated in such diverse cellular processes as proliferation, differentiation, and apoptosis. As each of these processes occurs during post-natal mammary gland morphogenesis, the expression and activity of NF- $\kappa$ B factors in the murine mammary gland were examined. Immunohistochemical and immunoblot analyses revealed expression of the p105/p50 and RelA subunits of NF- $\kappa$ B, as well as the major inhibitor, I $\kappa$ B $\alpha$ , in the mammary epithelium during pregnancy, lactation, and involution. Electrophoretic mobility shift assay (EMSA) demonstrated that DNA-binding complexes containing p50 and RelA were abundant during pregnancy and involution, but not during lactation. Activity of an NF- $\kappa$ B-dependent luciferase reporter in transgenic mice was highest during pregnancy, decreased to near undetectable levels during lactation, and was elevated during involution. This highly regulated pattern of activity was consistent with the modulated expression of p105/p50, RelA, and I $\kappa$ B $\alpha$ .



## Text

The Rel/NF- $\kappa$ B family of ubiquitous, inducible transcription factors consists of multiprotein complexes specialized for rapid cellular responses to a wide variety of growth factor, cytokine, and pathogenic stimuli (Verma et al., 1995). Although NF- $\kappa$ B activity has been well characterized in lymphoid tissues as a mediator of inflammatory and immune responses, several cell culture and *in vivo* models have implicated NF- $\kappa$ B in fundamental cellular processes, including proliferation and apoptosis (Verma et al., 1995; Attar et al., 1997; Ghosh, May, and Kopp, 1998). Each of these processes are critical during the highly dynamic morphogenesis of the mammary gland (Medina, 1996). Moreover, several recent studies have demonstrated aberrant expression of NF- $\kappa$ B factors and enhanced DNA binding activity in breast cancer cell lines and primary tumors (Dejardin et al., 1995; Nakashatri et al., 1997; Sovak et al., 1997; Sovak et al., 1999; Cogswell et al., 2000). Given these observations, the expression of several NF- $\kappa$ B/I $\kappa$ B family members over the course of post-natal mammary gland development was investigated. These family members included p105/p50, which acts as an NF- $\kappa$ B inhibitor in the 105 kDa isoform and as a transcriptional cofactor in the 50 kDa isoform, RelA, one of the major transcriptional activators in the NF- $\kappa$ B family, and I $\kappa$ B $\alpha$ , a direct inhibitor of NF- $\kappa$ B. In addition, the DNA-binding activity and *in vivo* transcription mediated by NF- $\kappa$ B throughout post-natal development was assessed.

Immunohistochemistry revealed expression of the p105/p50 and RelA subunits of NF- $\kappa$ B and the direct inhibitor, I $\kappa$ B $\alpha$ , in the epithelium of virgin, pregnant, lactating, and involuting mammary glands (Figure 1). Expression was observed in both the epithelium and in the surrounding stroma, with a more intense and dynamic pattern within the epithelium. At 16.5 days of pregnancy, when the mammary epithelium is proliferating and beginning to express milk proteins, RelA expression appeared to be upregulated as compared to levels in the virgin epithelium (Figure 1B). I $\kappa$ B $\alpha$  expression appeared to be decreased at 3 days of involution, when the mammary epithelium is regressing by apoptosis, compared to 5.5 day fully differentiated, lactating epithelium (Figure 1C). Specificity of the anti-p105/p50 antibody was confirmed by

probing sections from p105/p50-deficient virgin mammary glands. The ductal morphology and histology of p105/p50-deficient mammary glands appeared relatively normal, though a slight increase in lateral branching was observed in some samples (data not shown). Expression of p105/p50 was detected in wild-type virgin glands, while no staining was detected in glands collected from p105/p50-deficient virgin animals (Figure 1D). Specificity of anti-p105/p50 antisera, as well as antisera against RelA and I $\kappa$ B $\alpha$ , was also confirmed by peptide competition. The colorimetric signal in sections probed with antisera minus specific competitor was more intense than in control sections probed with primary antisera that was presorbed with specific competitor peptides, but not in sections probed with primary antisera that was presorbed with non-specific peptides (Figure 2). Higher magnification of p105/p50 and RelA stained sections revealed both nuclear and cytoplasmic localization of these subunits in epithelium during pregnancy and involution (Figure 3A).

In order to quantify p105/p50, RelA, and I $\kappa$ B $\alpha$  expression over the course of post-natal mammary gland morphogenesis, immunoblot analysis was performed on nuclear and cytoplasmic protein extracts from virgin, pregnant, lactating, and involuting mammary glands (Figure 3B). Enrichment of cytoplasmic and nuclear proteins within these extracts was confirmed by using antibodies directed against  $\beta$ -tubulin, a cytoplasmic protein, and SP1, a nuclear factor (Figure 3B). SP1 expression in nuclear extracts from involuting mammary samples was not observed, however, likely due to the cessation of much general transcription due to massive epithelial apoptosis during these stages. Moreover, the presence of intact proteins in these extracts and the relative uniformity of loading was confirmed by Ponceau S stain following transfer of SDS-PAGE fractionated proteins to polyvinylidene fluoride (PVDF) membranes (data not shown). I $\kappa$ B $\alpha$  functions as an inhibitor of NF- $\kappa$ B nuclear import primarily in the cytoplasm, and uniform I $\kappa$ B $\alpha$  expression was observed in cytoplasmic extracts from virgin, pregnant, and early lactating glands. Cytoplasmic expression of I $\kappa$ B $\alpha$  was observed to decrease in late lactation and involution. Cytoplasmic levels of latent p50 and RelA were also relatively uniform. Upon activation, NF- $\kappa$ B complexes are known to be transported to the nucleus, and an increase in nuclear RelA expression was observed

in pregnancy as compared to virgin samples, and again at 1 and 3 days involution as compared to 9.5 days lactation. Again, specificity of the p105/p50 antibody was confirmed by probing cytoplasmic extracts from wild-type virgin glands and from p105/p50-deficient glands with anti-p105/p50 antisera. Expression of p105/p50 was detected only in extracts from wild-type mammary glands, and uniform loading of proteins was confirmed by probing with an antibody against  $\beta$ -tubulin (Figure 3C). These data demonstrate a dynamic pattern of expression and intracellular localization for p105/p50, RelA, and I $\kappa$ B $\alpha$  in the mammary gland epithelium.

Electrophoretic mobility shift assays were performed on mammary gland nuclear extracts to assess the pattern of NF- $\kappa$ B DNA-binding activity during pregnancy, lactation, and involution. An oligonucleotide bearing an NF- $\kappa$ B consensus element from the human immunodeficiency virus-long terminal repeat (HIV-LTR) was used to detect nuclear NF- $\kappa$ B in these extracts. This enhancer binds a broad range of NF- $\kappa$ B homodimeric and heterodimeric complexes (Kretzshmar et al., 1992; Liu et al., 1992; Doerre et al., 1993). Two protein complexes capable of binding to this NF- $\kappa$ B consensus element were observed in mammary gland nuclear extracts. The relatively strong complex (Complex A) migrated with a higher mobility than the weaker complex (Complex B; Figure 4A). Specificity of the protein:DNA complex was demonstrated by the addition of excess unlabeled competitor oligonucleotides to 16.5 day pregnant, 5.5 day lactation, and 3 day post-wean samples (Figure 4A, lanes 3, 7, and 11). DNA binding activity was highest during pregnancy, decreased substantially through lactation, and then increased following cessation of nursing, as recently described by others (Clarkson and Watson, 1999). To provide a more detailed characterization of these DNA-binding complexes, antibodies recognizing specific family members were used for supershifts to identify specific NF- $\kappa$ B family members. In nuclear extracts prepared from mammary glands at 16.5 dpc, a supershift was produced by an antibody against p50 (Figure 4A, lane 5). RelA is also supershifted in nuclear extracts prepared from mammary glands at 16.5 dpc (Figure 4A, lane 6). RelA and p50 supershifts were also detected in nuclear extracts prepared from mammary glands at 3 dpw (Figure 4A, lanes 13 and 14). The specificity of the p105/p50 antibody used in supershift experiments was confirmed by incubating the antibody with

nuclear extracts from virgin p105/p50-deficient mammary glands. Nuclear extracts from p105/p50-deficient mice exhibited a decrease in the level of DNA-binding relative to extracts from wild-type mammary glands, although DNA-binding was still present (Figure 4B, lane 1 versus lane 5). No p50 supershift was observed in nuclear extracts prepared from virgin p105/p50-deficient mammary glands, although the anti-p50 antisera was able to produce a supershift in nuclear extracts prepared from wild-type virgin mammary glands (Figure 4B, lane 3 versus lane 7). No supershift was detected with the RelA antisera in p105/p50-deficient glands, though the more slowly migrating DNA-binding complex in p105/p50-deficient extracts appears to co-migrate with the more slowly migrating DNA-binding complex in wild-type virgins (Figure 4B, lane 4 versus lane 8). It is possible that this complex contains RelA but that the levels are below the detection limits in this assay (Figure 4B, lane 8). These data demonstrate the dynamic pattern DNA-binding activity of nuclear p50 and RelA containing complexes in the mammary gland during post-natal morphogenesis.

In order to assess the transactivational function of NF- $\kappa$ B *in vivo*, the activity of an NF- $\kappa$ B responsive reporter was monitored in the mammary glands of transgenic mice at various developmental stages. Transgenic mice bearing the human immunodeficiency virus long terminal repeat (HIV-LTR) driving the *Photinus* luciferase cDNA, termed HLLs, have been used to quantify both constitutive and induced NF- $\kappa$ B activity (Blackwell et al., in press). Moderate reporter activity was detected in virgin glands, compared to background levels in non-transgenic control glands (Figure 6). Luciferase activity increased consistently through mid-pregnancy (10.5 to 16.5 dpc), and then decreased in late pregnancy (18.5 dpc, Figure 5). NF- $\kappa$ B activity levels decreased dramatically between 3.5 and 5.5 days of lactation, matching the levels measured in non-transgenic control glands. When pups were removed after 9.5 days of lactation, to induce epithelial involution, an increase in NF- $\kappa$ B reporter activity was observed.

Immunohistochemistry and immunoblot analyses revealed specific expression of p105/p50, RelA, and I $\kappa$ B $\alpha$  members in the developing mammary epithelium. Nuclear p50 and RelA expression was upregulated during pregnancy and involution. In addition, EMSA and the activity

of an NF- $\kappa$ B inducible reporter in transgenic mice revealed elevated NF- $\kappa$ B activity during pregnancy and involution post-weaning. DNA-binding and transcriptional activity were dramatically reduced during lactation. These highly regulated and dynamic expression and activity patterns indicate that further studies are imperative to determine the specific functions of NF- $\kappa$ B in mammary epithelial development during pregnancy and involution.

## Materials and Methods

**Immunohistochemistry:** Paraformaldehyde-fixed, paraffin-embedded sections (5-7  $\mu\text{m}$ ) from number 3 thoracic glands of virgin, 16.5 day pregnant, 5.5 day lactating, and 3 day involuting (after forced wean at 9.5 days lactation) B6D2 mice were probed with primary antisera (rabbit anti-I $\kappa$ B $\alpha$ , SC-371, 1  $\mu\text{g}/\text{ml}$ ; rabbit anti-p105/p50, SC-114, 2  $\mu\text{g}/\text{ml}$ ; rabbit anti-RelA, SC-109, 5  $\mu\text{g}/\text{ml}$ ; Santa Cruz Biotechnology). For peptide competition assays, primary antisera were incubated with a 5 fold excess of specific peptides as per supplied protocol (Santa Cruz) prior to immunohistochemistry, or with a 5 fold excess of nonspecific peptides directed against p52 (SC-298P, Santa Cruz) as a negative control. Specific immunoreaction was detected using the Vectastain Elite ABC kit (Vector Laboratories) and 3,3',-Diaminobenzidine tetrahydrochloride (DAB) horseradish peroxidase substrate (Zymed Laboratories) according to manufacturers' protocols. Sections were counterstained with hematoxylin prior to dehydration, mounting, and photodocumentation.

**Immunoblot:** Nuclear and cytoplasmic extracts were prepared using number 4 inguinal mammary glands collected from virgin, 10.5 and 16.5 day pregnant, 5.5 and 9.5 day lactating, and 1,3 and 5 day post forced wean animals as described by Baldwin et al. (1991). Whole cell extracts from the FL5.12 pro-B cell line were used as a positive control for NF- $\kappa$ B/I $\kappa$ B expression, as progenitor B-cells are known to express these factors (Grumont and Genronidakis, 1994). Protein concentration was determined using a Lowry assay (Bio-Rad Protein Assay Kit). Protein extracts (15  $\mu\text{g}$  or 30  $\mu\text{g}$ ) were fractionated on 8 to 12% SDS-polyacrylamide gels and transferred to polyvinylidene fluoride (PVDF) membranes (Immobilon). The membranes were probed with primary antisera (rabbit anti-I $\kappa$ B $\alpha$  SC-371, 1  $\mu\text{g}/\text{ml}$ ; rabbit anti-p105/p50, NR1157, 1:1000, generously provided by Dr. Nancy Rice, National Cancer Institute ; rabbit anti-RelA, SC-109, 2  $\mu\text{g}/\text{ml}$ ; rabbit anti-SP1, SC-59, 1  $\mu\text{g}/\text{ml}$ ; mouse anti- $\beta$ -tubulin, N-357, 1:1000, Amersham) according to manufacturer's protocol (Santa Cruz Biotechnology). The specificity of the NR1157 antisera for immunoblot was confirmed previously by comparing expression in wild-type murine

thymic cell extracts versus extracts from p105/p50 deficient mice (Pereira et al., 1996). The membranes were then incubated with a horseradish peroxidase-conjugated goat anti-rabbit or goat anti-mouse secondary antisera (Southern Biotechnology Associates) diluted 1:3000. Specific proteins were visualized using Renaissance chemiluminescence reagents according to the manufacturer's instructions (NEN Life Sciences Products).

Electrophoretic Mobility Shift Assays (EMSAs): Conditions for nuclear extraction and NF- $\kappa$ B EMSA have been detailed elsewhere (Baldwin et al., 1991). Oligonucleotides encoding an NF- $\kappa$ B consensus element from the HIV-LTR were used as probes (Stratagene Gelshift Assay Kit; plus strand 5'- GAT CGA GGG GAC TTT CCC TAG C- 3'). Briefly, 10  $\mu$ g of nuclear extracts from number 4 inguinal mammary glands of virgin, 10.5 and 16.5 day pregnant, 5.5 and 9.5 day lactating, and 1, 3, and 5 days involuting B6D2 mice were incubated in binding buffer containing 1  $\mu$ g poly(dI-dC) (Boehringer-Mannheim) and, where appropriate, 2  $\mu$ g specific antiserum (anti-RelA, SC-109X, anti-p50, SC-114X; Santa Cruz Biotechnology). Probe (30,000 cpm) was added to each sample, and the samples were incubated for 25 minutes at ambient temperature. Samples were fractionated on a 4% polyacrylamide gel in 0.5X TBE for 2 hours at 150V. The gel was dried and exposed to autoradiographic film (Kodak) for 1-2 days at -80°C.

p50 Deficient Mice: Mice deficient for p105/p50 were obtained from the Jackson Laboratory (B6.129P - Nfkb1[tm1Bal]). The genotype was confirmed by polymerase chain reaction (PCR) analysis as per supplier's protocol using the following primers: common primer 5' - GCA AAC CTG GGA ATA CTT CAT GTG ACT AAG - 3', wild-type primer 5' - ATA GGC AAG GTC AGA ATG CAC CAG AAG TCC - 3', and null primer 5' - AAA TGT GTC AGT TTC ATA GCC TGA AGA ACG - 3'.

Generation of HIV-LTR-luciferase (HLL) transgenic animals and luciferase assay: The HIV-LTR-luciferase transgene consists of a 680 bp HIV-LTR XhoI/HindIII fragment from an HIV-LTR lacZ plasmid derived from the pU3R-III HIV-LTR-CAT plasmid (Sodroski et al., 1985; Nabel et al., 1987) driving luciferase from the pGL2-Basic vector (Promega). Transgenic animals were generated by pronuclear injection (Vanderbilt University Cancer Center Transgenic Core Facility),

and two transgenic founders were identified and used to generate two expressing lines (20 and 27). Presence of the luciferase transgene was assessed by Southern blot and/or PCR analysis of genomic DNA from tail biopsy, and confirmed by constitutive luciferase activity in the brain (Blackwell et al., in press). Basal expression of the luciferase reporter, as well as inducible expression upon treatment with bacterial lipopolysaccharide (LPS) was used to validate these mice as a model system for NF- $\kappa$ B-mediated transcriptional activity *in vivo* (Blackwell et al., in press).

Total cell extracts were prepared in passive lysis buffer (Analytical Bioluminescence) from number 4 inguinal mammary glands of HLL +/- transgenic mice and wild-type littermates at the following stages: virgin; 10.5, 12.5, 14.5, 16.5, and 18.5 days pregnancy; 1.5, 3.5, 5.5, 7.5, and 9.5 days lactation; 1, 3, and 5 days involution. Luciferase activity was measured according to the manufacturer's instructions (Analytical Bioluminescence). Protein concentrations in the extracts were determined by Lowry assay (Bio-Rad). All values are presented as relative light units (RLUs)/ $\mu$ g protein.



cell extracts versus extracts from p105/p50 deficient mice (Pereira et al., 1996). The lanes were then incubated with a horseradish peroxidase-conjugated goat anti-rabbit or goat mouse secondary antisera (Southern Biotechnology Associates) diluted 1:3000. Specific bands were visualized using Renaissance chemiluminescence reagents according to the manufacturer's instructions (NEN Life Sciences Products).

Electrophoretic Mobility Shift Assays (EMSAs): Conditions for nuclear extraction and NF- $\kappa$ B assays have been detailed elsewhere (Baldwin et al., 1991). Oligonucleotides encoding an NF- $\kappa$ B consensus element from the HIV-LTR were used as probes (Stratagene Gelshift Assay Kit; plus 5'-GAT CGA GGG GAC TTT CCC TAG C-3'). Briefly, 10  $\mu$ g of nuclear extracts from 4 inguinal mammary glands of virgin, 10.5 and 16.5 day pregnant, 5.5 and 9.5 day nursing, and 1, 3, and 5 days involuting B6D2 mice were incubated in binding buffer containing 1  $\mu$ g (dI-dC) (Boehringer-Mannheim) and, where appropriate, 2  $\mu$ g specific antiserum (anti-SC-109X, anti-p50, SC-114X; Santa Cruz Biotechnology). Probe (30,000 cpm) was added to each sample, and the samples were incubated for 25 minutes at ambient temperature. Gels were fractionated on a 4% polyacrylamide gel in 0.5X TBE for 2 hours at 150V. The gel was stained and exposed to autoradiographic film (Kodak) for 1-2 days at -80°C.

Deficient Mice: Mice deficient for p105/p50 were obtained from the Jackson Laboratory (29P - Nfkb1[tm1Bal]). The genotype was confirmed by polymerase chain reaction (PCR) analysis as per supplier's protocol using the following primers: common primer 5' - GCA AAC GGA ATA CTT CAT GTG ACT AAG - 3', wild-type primer 5' - ATA GGC AAG GTC ATG CAC CAG AAG TCC - 3', and null primer 5' - AAA TGT GTC AGT TTC ATA GCC AGA ACG - 3'.

Generation of HIV-LTR-luciferase (HLL) transgenic animals and luciferase assay: The HIV-LTR-luciferase transgene consists of a 680 bp HIV-LTR XhoI/HindIII fragment from an HIV-LTR lacZ reporter construct derived from the pU3R-III HIV-LTR-CAT plasmid (Sodroski et al., 1985; Nabel et al., 1987) driving luciferase from the pGL2-Basic vector (Promega). Transgenic animals were generated by pronuclear injection (Vanderbilt University Cancer Center Transgenic Core Facility),

and two transgenic founders were identified and used to generate two expressing lines (20 and 27). Presence of the luciferase transgene was assessed by Southern blot and/or PCR analysis of genomic DNA from tail biopsy, and confirmed by constitutive luciferase activity in the brain (Blackwell et al., in press). Basal expression of the luciferase reporter, as well as inducible expression upon treatment with bacterial lipopolysaccharide (LPS) was used to validate these mice as a model system for NF- $\kappa$ B-mediated transcriptional activity *in vivo* (Blackwell et al., in press).

Total cell extracts were prepared in passive lysis buffer (Analytical Bioluminescence) from number 4 inguinal mammary glands of HLL +/- transgenic mice and wild-type littermates at the following stages: virgin; 10.5, 12.5, 14.5, 16.5, and 18.5 days pregnancy; 1.5, 3.5, 5.5, 7.5, and 9.5 days lactation; 1, 3, and 5 days involution. Luciferase activity was measured according to the manufacturer's instructions (Analytical Bioluminescence). Protein concentrations in the extracts were determined by Lowry assay (Bio-Rad). All values are presented as relative light units (RLUs)/ $\mu$ g protein.

## Acknowledgments

We thank the Vanderbilt University Department of Histopathology and the Vanderbilt University Skin Disease Research Center for preparation of mouse mammary gland sections. We would also like to thank Annapurna Venkatakrishnan from the laboratory of Dr. Timothy S. Blackwell at Vanderbilt University for providing labeled NF- $\kappa$ B probes for EMSA. We also thank Dr. Nancy Rice at the National Cancer Institute for providing the NR1157 anti-p50 antisera. Special thanks to members of the Kerr lab and to Drs. Carlos Arteaga, Lynn Matrisian, Harold Moses, Virginia Shepherd, and David Threadgill at Vanderbilt University for helpful discussion and/or review of the manuscript. Dana M. Brantley is supported by a Department of Defense Breast Cancer Research Predoctoral Fellowship (DAMD17-97-1-7057). This work was supported by NIH grant R01GM51249, a Center grant from the National Cancer Institute (CA 68485), American Cancer Society Grant IRG-25-38, and by funding for Breast Cancer Research Project #9838 from the Susan G. Komen Breast Cancer Foundation.

## Figure Legends

Figure 1: NF- $\kappa$ B and I $\kappa$ B $\alpha$  are expressed in the mammary gland epithelium over the course of post-natal morphogenesis. Histological sections were prepared from virgin, 16.5 day pregnant (16.5 dpc), 5.5 day lactating (5.5 lac), and 3 days post-wean involuting (3 dpw) number 3 thoracic glands. These sections were probed with primary antisera for p105/p50 (A), RelA (B), or I $\kappa$ B $\alpha$  (C). Epithelial expression of RelA increased at 16.5 days of pregnancy (B), whereas expression of I $\kappa$ B $\alpha$  decreased at 3 days post-wean involution. Histological sections from wild-type and p105/p50-deficient mammary glands were probed with antisera against p105/p50 to confirm the specificity of the antisera. Expression of p105/p50 was detected in sections from wild-type virgin animals but not in sections from p105/p50-deficient virgin animals (D). Data presented are a representative presentation of results from 3 to 6 independent samples per developmental stage. Scale bar = 50  $\mu$ m.

Figure 2: Specific expression of NF- $\kappa$ B and I $\kappa$ B $\alpha$  in the mammary gland. Histological sections were prepared from number 3 thoracic glands collected from 3 days post-wean involuting (3 dpw) mice as both NF- $\kappa$ B factors and I $\kappa$ B $\alpha$  are expressed at detectable levels during this stage of development. These sections were probed with primary antisera for p105/p50 (A), RelA (B), or I $\kappa$ B $\alpha$  (C), with or without 5 fold excess of specific competitor peptides or nonspecific peptides against p52. The specific peptides abrogated the signal (left hand panel versus middle), whereas the nonspecific peptides did not (left hand panel versus right), demonstrating the specificity of these antisera for NF- $\kappa$ B factors and I $\kappa$ B $\alpha$ . Data presented are a representation of results from 3 independent samples. Scale bar = 50  $\mu$ m.

Figure 3: NF- $\kappa$ B factors are localized to both the nuclei and cytoplasm of mammary epithelium during pregnancy and involution. Higher magnification of 16.5 dpc and 3 dpw sections probed for p105/p50 and RelA expression revealed nuclear localization in several epithelial cells (A, top row) in addition to the cytoplasmic localization in adjacent cells. Scale bar = 10  $\mu$ m. Western blot analysis of cytoplasmic and nuclear protein extracts (15  $\mu$ g) from mammary glands or FL5.12 pro-B cells as a negative control were subjected to immunoblot analysis for expression of I $\kappa$ B $\alpha$ , p105/p50, and RelA (B) or for  $\beta$ -tubulin and SP1 as cytoplasmic and nuclear markers respectively. I $\kappa$ B $\alpha$  expression in the cytoplasm was uniform until 9.5 days lactation (9.5 lac), when levels decreased during through 1, 3 and 5 days involution (1, 3, 5 dpw). Expression of p50 and RelA was relatively uniform in the cytoplasm. Nuclear RelA expression increased in pregnancy and involution. The specificity of the p105/p50 antisera was confirmed by immunoblot analysis using cytoplasmic extracts (30  $\mu$ g) from wild-type or p105/p50-deficient virgin mammary glands (C). Expression of p105/p50 was detected in cytoplasmic extracts prepared from wild-type mammary glands, but not in extracts prepared from p105/p50-deficient glands. Antisera against  $\beta$ -tubulin was used to confirm that equal quantities of protein were loaded for each sample (C). Data presented are a representation of results from 3 to 6 independent samples per developmental stage.

Figure 4: DNA binding activity of NF- $\kappa$ B in mammary gland nuclear extracts is highest during pregnancy and involution. Nuclear extracts were prepared from virgin, pregnant (10.5 dpc and 18.5 dpc), lactating (5.5 lac and 9.5 lac), and involuting (1 dpw, 3 dpw, and 5 dpw) mammary glands. Nuclear extracts (10  $\mu$ g) were incubated with an end-labeled oligonucleotide containing an NF- $\kappa$ B enhancer element from the HIV-LTR. Two protein complexes capable of binding this oligonucleotide were present in virgin, pregnant, and involuting mammary samples, a strong complex (Complex A) migrating below a relatively weaker complex (Complex B; A). The specificity of these complexes was confirmed by competition with excess, unlabeled oligonucleotide (A; lanes 3, 7, and 11). Supershifts for p50 and RelA confirmed the presence of the NF- $\kappa$ B components in the complex (A; lanes 5, 6, 13, and 14). In order to confirm the specificity of the antisera used to supershift p50, nuclear extracts prepared from virgin p105/p50-deficient mice were probed with anti-p50 antisera. DNA-binding in p105/p50-deficient nuclear extracts (10  $\mu$ g) was apparent, although reduced compared to that in wild-type nuclear extracts (B; lane 1 versus lane 5). The antisera used to detect p50 in EMSA experiments was specific, as a supershift was detected in wild-type but not p105/p50-deficient nuclear extracts (B; lane 3 versus lane 7). Data presented are a representation of 3 to 6 independent samples per developmental stage.

Figure 5: *In vivo* transcriptional activity of NF- $\kappa$ B in the mammary glands of transgenic reporter mice is highest during pregnancy and involution. Protein extracts were prepared from virgin, pregnant (10.5 dpc, 12.5 dpc, 14.5 dpc, 16.5 dpc, and 18.5 dpc), lactating (1.5 lac, 3.5 lac, 5.5 lac, 7.5 lac, and 9.5 lac), and involuting (1 dpw, 3 dpw, and 5 dpw) number 4 inguinal mammary glands from transgenic mice expressing luciferase under the regulation of the HIV-LTR or from non-transgenic animals at 10.5 days of pregnancy. Luciferase activity (relative light units, RLUs) was assessed in 20  $\mu$ l of protein extract. Data were normalized based on protein concentration and are the average of 3 independent samples per developmental stage with standard error of the mean.



## ferences

- Far, R. M., Caamano, J., Carrasco, D., Violetta, I., Ishikawa, H., Ryseck, R.-P., Weih, F., and Bravo, R. 1997. Genetic approaches to study Rel/NF- $\kappa$ B/I $\kappa$ B function in mice. *Cancer Biol.* 93-101.
- Baldwin, A. S., Jr., Azizkhan, J. C., Jensen, D. E., Beg, A. A., and Coodly, L. R. 1991. Reduction of NF- $\kappa$ B DNA-binding activity during the G0-to-G1 transition in mouse fibroblasts. *Mol. Cell. Biol.* 11, 4943-51.
- Blackwell, T. S., Yull, F. E., Chen, C.-L., Venkatakrishna, A., Blackwell, T. R., Hicks, D. J., Lancaster, L. H., Christman, J. W., and Kerr, L. D. 2000. Multi-organ NF- $\kappa$ B activation in a transgenic mouse model of systemic inflammation. *Am. J. Respir. Crit. Care Med.* In press.
- Markson, R. W. E., and Watson, C. J. 1999. NF- $\kappa$ B and apoptosis in mammary epithelial cells. *J. Mammary Gland Biol. Neo.* 4, 165-175.
- Ngswell, P. C., Guttridge, D. C., Funkhouser, W. K., and Baldwin, A. S., Jr. 2000. Selective inactivation of NF- $\kappa$ B subunits in human breast cancer: potential roles for NF- $\kappa$ B2/p52 and for Bcl-2. *Oncogene.* 19, 1123-31.
- Quardin, E., Bonizzi, G., Bellahcene, A., Castronovo, V., Merville, M. P., and Bours, V. 2000. Highly-expressed p100/p52 (NFKB2) sequesters other NF- $\kappa$ B-related proteins in the cytoplasm of human breast cancer cells. *Oncogene.* 11, 1835-41.
- Perre, S., Sista, P., Sun, S. C., Ballard, D. W., and Greene, W. C. 1993. The c-rel proto-oncogene product represses NF- $\kappa$ B p65-mediated transcriptional activation of the long terminal repeat of type 1 human immunodeficiency virus. *Proc. Natl. Acad. Sci. U S A* 90, 1023-27.
- Rossh, S., May, M. J., and Kopp, E. B. 1998. NF- $\kappa$ B and rel proteins: evolutionarily conserved mediators of immune responses. *Annu. Rev. Immunol.* 16, 225-60.
- Schmont, R. J. and Gerondakis, S. 1994. The subunit composition of NF-kappa B complexes changes during B-cell development. *Cell Growth Differ.* 5, 1321-31.
- Stettschmar, M., Meisterernst, M., Scheidereit, C., Li, G., and Roeder, R. G. 1992. Transcriptional regulation of the HIV-1 promoter by NF- $\kappa$ B *in vitro*. *Genes Dev.* 6, 761-74.
- Thompson, J., Perkins, N. D., Schmid, R. M., and Nabel, G. J. 1992. Specific NF- $\kappa$ B subunits act in concert with Tat to stimulate human immunodeficiency virus type 1 transcription. *J. Virol.* 66, 13-7.
- Udina, D. 1996. The mammary gland: a unique organ for the study of development and carcinogenesis. *J. Mammary Gland Biol. Neo.* 1, 5-19.
- Waghshatri, H., Bhat-Nakshatri, P., Martin, D. A., Goulet, R. J., Jr., and Sledge, G. W., Jr. 1997. Constitutive activation of NF- $\kappa$ B during progression of breast cancer to hormone-independent growth. *Mol. Cell Biol.* 17, 3629-39.
- Wiel, G., and Baltimore, D. 1987. An inducible transcription factor activates expression of human immunodeficiency virus in T cells. *Nature* 326, 711-3.

ira, D. S., Kushner, D. B., Ricciardi, R. P., and Graham, F. L. 1996. Testing NF- $\kappa$ B1-p50 body specificity using knockout mice. *Oncogene* 13, 445-446.

roski, J., Rosen, C., Wong-Staal, F., Salahuddin, S. Z., Popovic, M., Arya, S., Gallo, R. and Haseltine, W. A. 1985. Trans-acting transcriptional regulation of human T-cell leukemia virus type III long terminal repeat. *Science* 227, 171-3.

ak, M. A., Bellas, R. E., Kim, D. W., Zanieski, G. J., Rogers, A. E., Traish, A. M., and Sonenshein, G. E. 1997. Aberrant nuclear factor- $\kappa$ B/Rel expression and the pathogenesis of breast cancer. *J. Clin. Invest.* 100, 2952-60.

ak, M. A., Arsura, M., Zanieski, G., Kavanagh, K. T., and Sonenshein, G. E. 1999. The mitogenic effects of transforming growth factor  $\beta$ 1 on breast cancer cell proliferation are mediated through regulation of aberrant nuclear factor- $\kappa$ B/rel expression. *Cell Growth and Diff.* 10, 537-

na, I. M., Stevenson, J. K., Schwarz, E. M., Van Antwerp, D., and Miyamoto, S. 1995. NF- $\kappa$ B/I $\kappa$ B family: intimate tales of association and dissociation. *Genes Dev.* 9, 2723-35.